

## Some results on generalized $(k, \mu)$ -space forms

Shanmukha B<sup>1</sup>, Venkatesha<sup>1</sup> and Vishunuvardhana S. V.<sup>2</sup>

<sup>1</sup>Department of Mathematics, Kuvempu University, Shankaraghatta, Shimoga, Karnataka, India

<sup>2</sup>Department of Mathematics, Gitam School of technology, Gitam (Deemed to be University), Bangalore, Karnataka, India

Received: 18 December 2017, Accepted: 25 March 2018

Published online: 22 July 2018.

**Abstract:** In this paper we have studied Ricci symmetric and Ricci pseudosymmetric generalized  $(k, \mu)$ -space forms and generalized  $(k, \mu)$ -space forms with quasi umblical hypersurface and  $\tau$ -flat curvature tensor.

**Keywords:** Generalized  $(k, \mu)$ -space form, Ricci symmetric, Ricci pseudosymmetric, quasi umblical hypersurface,  $\tau$ -curvature tensor.

### 1 Introduction

An almost contact metric manifold  $(M, g)$  is a Riemannian manifold with a tensor field  $\phi$  of type  $(1, 1)$ , a vector field  $\xi$ , a 1-form  $\eta$  on  $M$  satisfying [5, 6]

$$\phi^2 = -I + \eta \circ \xi, \quad \eta(\xi) = 1, \quad \eta \circ \phi = 0, \quad \phi\xi = 0, \quad (1)$$

$$g(\phi X, \phi Y) = g(X, Y) - \eta(X)\eta(Y), \quad (2)$$

$$g(\phi X, Y) = -g(X, \phi Y), \quad g(\phi X, X) = 0, \quad g(X, \xi) = \eta(X), \quad (3)$$

for all vector fields  $X, Y$  on  $M$ . We know that a real space form is a Riemannian manifold having constant sectional curvature and a complex space form is a Kaehlerian manifold  $(M, J, g)$  with constant holomorphic sectional curvature  $c$ .

Generalized Sasakian space forms were studied extensively in [1, 2, 3, 15, 18, 19, 25]. An almost contact metric manifold  $(M, \phi, \xi, \eta, g)$  is a generalized  $(k, \mu)$  space form if there exists differential functions  $f_1, f_2, \dots, f_6$  on  $M^{2n+1}(f_1, \dots, f_6)$ , whose curvature tensor  $R$  is given by [8, 9]

$$R = f_1 R_1 + f_2 R_2 + f_3 R_3 + f_4 R_4 + f_5 R_5 + f_6 R_6, \quad (4)$$

where  $R_1, R_2, R_3, R_4, R_5$  and  $R_6$  are given by;

$$R_1(X, Y)Z = \{g(Y, Z)X - g(X, Z)Y\},$$

$$R_2(X, Y)Z = \{g(X, \phi Z)\phi Y - g(Y, \phi Z)\phi X + 2g(X, \phi Y)\phi Z\},$$

$$R_3(X, Y)Z = \{\eta(X)\eta(Z)Y - \eta(Y)\eta(Z)X + g(X, Z)\eta(Y)\xi - g(Y, Z)\eta(X)\xi\}$$

$$R_4(X, Y)Z = \{g(Y, Z)hX - g(X, Z)hY + g(hY, Z)X - g(hX, Z)Y\},$$

$$R_5(X, Y)Z = g(hY, Z)hX - g(hX, Z)hY + g(\phi hX, Z)\phi hY - g(\phi hY, Z)\phi hX,$$

$$R_6(X, Y)Z = \eta(X)\eta(Z)hY - \eta(Y)\eta(Z)hX + g(hX, Z)\eta(Y)\xi - g(hY, Z)\eta(X)\xi,$$

Here  $h$  is defined by  $2h = L_\xi \phi$  is a symmetric tensor and satisfies the following conditions

$$h\xi = 0, \quad h\phi = -\phi h, \quad \text{tr}(h) = 0, \quad \eta \circ h = 0, \quad (5)$$

and where  $L$  is the usual Lie derivative. In particular if  $f_4 = f_5 = f_6 = 0$ , then generalized  $(k, \mu)$ -space form  $M^{2n+1}(f_1, \dots, f_6)$  reduces to generalized Sasakian space forms. Also in [17] it was proved that  $(k, \mu)$ -space forms are natural examples of generalized  $(k, \mu)$ -space forms for constant functions  $f_1 = \frac{c+3}{4}$ ,  $f_2 = \frac{c-1}{4}$ ,  $f_3 = \frac{c+3}{4} - k$ ,  $f_4 = 1$ ,  $f_5 = \frac{1}{2}$ ,  $f_6 = 1 - \mu$ .

In a generalized  $(k, \mu)$  space forms, the following relations hold [9];

$$S(X, Y) = (2nf_1 + 3f_2 - f_3)g(X, Y) - (3f_2 + (2n - 1)f_3)\eta(X)\eta(Y) + ((2n - 1)f_4 - f_6)g(hX, Y), \quad (6)$$

$$R(X, Y)\xi = (f_1 - f_3)\{\eta(Y)X - \eta(X)\eta(Y)\} + (f_4 - f_6)\{\eta(Y)hX - \eta(X)hY\}, \quad (7)$$

$$S(\phi X, \phi Y) = S(X, Y) - 2n(f_1 - f_3)\eta(X)\eta(Y). \quad (8)$$

**Definition 1.** A Riemannian manifold  $M$  is said to be

- (I) Einstein manifold if  $S(X, Y) = \lambda_1 g(X, Y)$ ,
- (II)  $\eta$ -Einstein manifold if  $S(X, Y) = \lambda_1 g(X, Y) + \lambda_2 \eta(X)\eta(Y)$ ,
- (III) Special type of  $\eta$ -Einstein manifold if  $S(X, Y) = \lambda_1 \eta(X)\eta(Y)$ ,

where  $S$  is the Ricci tensor and  $\lambda_1$  and  $\lambda_2$  are constants.

In a generalized quasi-Einstein manifold the Ricci tensor  $S$  is given by [13]

$$S(X, Y) = \alpha g(X, Y) + \beta A(X)A(Y) + \gamma B(X)B(Y), \quad (9)$$

Here  $\alpha, \beta, \gamma$  are non zero scalars and  $A, B$  are non-zero 1-forms which are defined by

$$g(X, U) = A(X) \text{ and } g(X, V) = B(X),$$

where  $U$  and  $V$  are two orthogonal vectors. If  $\gamma = 0$ , then the manifold reduces to a quasi Einstein manifold.

## 2 Main results

**Theorem 1.** Let  $M$  be quasi-umbilical hypersurface of a generalized  $(k, \mu)$ -space form is a generalized quasi-Einstein hypersurface if and only if  $f_6 = (2n - 1)f_4$ .

*Proof.* We know that hypersurface of  $(M^{2n+1}, \tilde{g})$  is  $(M^{2n}, g)$ . If  $A$  is the  $(1, 1)$ -tensor corresponding to the normal valued second fundamental tensor  $H$ , then we have [11]

$$g(A_\rho(X), Y) = \tilde{g}(H(X, Y), \rho), \quad (10)$$

where  $\rho$  is a unit normal vector field and  $X, Y$  are tangent vector fields. Let  $H_\rho$  be the symmetric  $(0, 2)$  tensor corresponding to  $A_\rho$  in the hypersurface, defined by

$$g(A_\rho(X), Y) = H_\rho(X, Y). \quad (11)$$

A hypersurface of a Riemannian manifold  $(M^{2n+1}, g)$  is called quasi-umbilical if its second fundamental tensor has the form [11]

$$H_\rho(X, Y) = \alpha g(X, Y) + \beta \omega(X)\omega(Y), \quad (12)$$

where  $\omega$  is a 1-form and  $\alpha, \beta$  are scalars. Suppose  $\alpha = 0$  (resp.  $\beta = 0$  or  $\alpha = \beta = 0$ ) holds, then it is called cylindrical (resp. umbilical or geodesic). Now from (10), (11) and (12) we obtain

$$g(H(X, Y), \rho) = \alpha g(X, Y) \tilde{g}(\rho, \rho) + \beta \omega(X) \omega(Y) \tilde{g}(\rho, \rho),$$

which implies that

$$H(X, Y) = \alpha g(X, Y) \rho + \beta \omega(X) \omega(Y) \rho. \tag{13}$$

The Gauss equation tangent to the hypersurface is given by [11]

$$R(X, Y, Z, W) = \tilde{R}(X, Y, Z, W) - g(H(X, W), H(Y, Z)) + g(H(X, Z), H(Y, W)), \tag{14}$$

where  $\tilde{R}(X, Y, Z, W) = \tilde{g}(R(X, Y)Z, W)$  and  $R(X, Y, Z, W) = g(R(X, Y)Z, W)$ .

Let us consider quasi umbilical hypersurface of generalized  $(k, \mu)$ -space forms. Then from (13) and (14) we have

$$R(X, Y, Z, W) = \tilde{R}(X, Y, Z, W) - g([\alpha g(X, W) \rho + \beta \omega(X) \omega(W) \rho], [\alpha g(Y, Z) \rho + \beta \omega(Y) \omega(Z) \rho]) + g([\alpha g(Y, W) \rho + \beta \omega(Y) \omega(W) \rho], [\alpha g(X, Z) \rho + \beta \omega(X) \omega(Z) \rho]). \tag{15}$$

Using (4) in (15) and then contracting over  $X$  and  $W$ , we get

$$S(Y, Z) = (2nf_1 + 3f_2 - f_3 + 2n\alpha^2 + \alpha^2)g(Y, Z) - (3f_2 + (2n - 1)f_3)\eta(Y)\eta(Z) + ((2n - 1)f_4 - f_6)g(hY, Z) + (2n - 1)\alpha\beta\omega(Y)\omega(Z). \tag{16}$$

This complete the proof of the theorem

**Theorem 2.** A generalized  $(k, \mu)$ -space form  $M^{2n+1}(f_1, \dots, f_6)$  satisfying  $(S(X, \xi) \cdot R)(Y, Z)W = 0$  is an  $\eta$ -Einstein manifold.

*Proof.* Consider a generalized  $(k, \mu)$ -space form  $(n > 1)$  satisfying  $(S(X, \xi) \cdot R)(Y, Z)W = 0$ , then we have

$$\begin{aligned} 0 &= (S(X, \xi) \cdot R)(Y, Z)W = (X \wedge_s \xi)R(Y, Z)W \\ &= ((X \wedge_s \xi) \cdot R)(Y, Z)W + R((X \wedge_s \xi)Y, Z)W \\ &\quad + R(Y, (X \wedge_s \xi)Z)W + R(Y, Z)(X \wedge_s \xi)W, \end{aligned} \tag{17}$$

where  $(X \wedge_s Y)$  is an endomorphism and is defined by

$$(X \wedge_s Y)Z = S(Y, Z)X - S(X, Z)Y. \tag{18}$$

Using (7) and (18) in (17) and taking inner product with  $\xi$  to the resulting equation, then we have

$$\begin{aligned} &2n(f_1 - f_3)\{\eta(R(Y, Z)W)\eta(X) + \eta(Y)\eta(R(X, Z)W) + \eta(Z)\eta(R(Y, X)W) + \eta(W)\eta(R(Y, Z)X)\} \\ &- \{S(X, R(Y, Z)W) + S(X, Y)\eta(R(\xi, Z)W) + S(X, Z)\eta(R(Y, \xi)W) + S(X, W)\eta(R(Y, Z)\xi)\} = 0. \end{aligned}$$

Setting  $Y = W = \xi$  in the above equation, we get

$$\begin{aligned} &(f_1 - f_3)\{2n(f_1 - f_3)\eta(X)\eta(Z) - S(X, Z)\} + 2n(f_1 - f_3)^2\{g(X, Z) - \eta(Z)\eta(X) \\ &+ (f_4 - f_6)2n(f_1 - f_3)g(hZ, X)\} = 0. \end{aligned} \tag{19}$$

By virtue of (6) in (19), we have

$$S(X, Z) = Ag(X, Z) + B\eta(X)\eta(Z),$$

where

$$A = -\left[\frac{((2n-1)f_4 - f_6)2n(f_1 - f_3) - (2nf_1 + 3f_2 - f_3)}{2n(f_1 - f_3)((2n-1)f_4 - f_6) + (f_4 - f_6)2n}\right]$$

and

$$B = -\left[\frac{-4n(f_1 - f_3) + (3f_2 + (2n-1)f_3)}{2n(f_1 - f_3)((2n-1)f_4 - f_6) + (f_4 - f_6)2n}\right].$$

Hence the proof.

**Definition 2.A** *generalized  $(k, \mu)$ -space forms is said to be Ricci symmetric if it satisfies*

$$(\nabla_W S)(X, Y) = 0$$

for all  $X, Y$  are orthogonal to  $\xi$ .

**Theorem 3.** *Let  $M$  be a 3-dimensional generalized  $(k, \mu)$ -space form has constant  $\phi$ -sectional curvature with constants  $(f_1 - f_3)$  and  $(f_4 - f_6)$ , then the following are equivalent:*

- (1) Ricci symmetric with  $(f_4 - f_6) \neq 0$ ,
- (2) Tensor  $h$  is parallel.

*Proof.* Differentiating (6) with respect to  $W$ , we get

$$\begin{aligned} (\nabla_W S)(X, Y) &= \{2nd(f_1 - f_3)(W) + 3df_2(W) + (2n-1)df_3(W)\}g(X, Y) - \{3df_2(W) + (2n-1)df_3(W)\}\eta(X)\eta(Y) \\ &\quad + \{2(n-1)df_4(W) + d(f_4 - f_6)(W)\}g(hX, Y) - (3f_2 + (2n-1)f_3)\{(\nabla_W \eta)(X)\eta(Y) \\ &\quad + (\nabla_W \eta)(Y)\eta(X)\} + \{2(n-1)f_4 + (f_4 - f_6)\}g((\nabla_W h)X, Y). \end{aligned} \quad (20)$$

If  $X$  and  $Y$  are orthogonal to  $\xi$ , then we have from (20)

$$\begin{aligned} (\nabla_W S)(X, Y) &= \{2nd(f_1 - f_3)(W) + 3df_2(W) + (2n-1)df_3(W)\}g(X, Y) + \{2(n-1)df_4(W) \\ &\quad + d(f_4 - f_6)(W)\}g(hX, Y) + \{2(n-1)f_4 + (f_4 - f_6)\}g((\nabla_W h)X, Y). \end{aligned} \quad (21)$$

Suppose  $(f_1 - f_3)$  and  $(f_4 - f_6)$  are non-zero constants, then (21) becomes

$$(\nabla_W S)(X, Y) = \{3df_2(W) + (2n-1)df_3(W)\}g(X, Y) + \{2(n-1)df_4(W)\}g(hX, Y) + \{(2n-1)f_4 - f_6\}g((\nabla_W h)X, Y). \quad (22)$$

For  $n = 1$  (i.e. for three-dimensional case) in (22), we have

$$(\nabla_W S)(X, Y) = d(3f_2 + f_3)(W)g(X, Y) + (f_4 - f_6)g((\nabla_W h)X, Y). \quad (23)$$

Let the  $\phi$ -sectional curvature  $(3f_2 + f_3)$  ( see [9]) of generalized  $(k, \mu)$ -space form be constant. Then (23) gives

$$(\nabla_W S)(X, Y) = (f_4 - f_6)g((\nabla_W h)X, Y). \quad (24)$$

Hence the proof, moreover.

**Theorem 4.** *Let  $(2n+1)$ -dimensional  $(k, \mu)$ -space form with  $X, Y \in \xi^\perp$ , then the following are equivalent:*

- (I) Ricci symmetric with  $\{(2n-1)f_4 - f_6\} \neq 0$ ,
- (II) Tensor  $h$  is parallel.

*Proof.* Consider the functions  $f_1, \dots, f_6$  in (4) are constants, then generalized  $(k, \mu)$ -space form reduces to the  $(k, \mu)$ -space form. Now suppose that the functions  $f_1 \dots f_6$  are constants. Then by taking covariant differentiation of (6) with respect to  $W$ , we get

$$(\nabla_W S)(X, Y) = -(3f_2 + (2n - 1)f_3)\{(\nabla_W \eta)(X)\eta(Y) + \eta(X)(\nabla_W \eta)(Y)\} + \{(2n - 1)f_4 - f_6\}g((\nabla_W h)X, Y). \quad (25)$$

If  $X, Y$  are orthogonal to  $\xi$ , then (25) yields

$$(\nabla_W S)(X, Y) = \{(2n - 1)f_4 - f_6\}g((\nabla_W h)X, Y).$$

This completes the proof of the theorem.

When generalizing the spaces of constant curvature, a locally symmetric spaces were introduced by Cartan [7]. All locally symmetric space satisfies  $R \cdot R = 0$ , where the first  $R$  represents the curvature operator which acts as a derivation and the second  $R$  represents the Riemannian curvature tensor. Manifolds satisfying the condition  $R \cdot R = 0$  are called semisymmetric manifolds and were classified by Szabo [24]. The condition of semisymmetry was weakened by Deszcz as pseudosymmetry which are characterized by the condition  $R \cdot R = LQ(g, R)$ , here by  $L$  is a real function and  $Q(g, R)$  is the Tachibana tensor.

**Definition 3.** A Riemannian manifold  $M$  is said to be pseudosymmetric, in the sense of Deszcz [14] if

$$(R(X, Y) \cdot R)(Z, U)V = L_R\{((X \wedge Y) \cdot R)(Z, U)V\}, \quad (26)$$

holds. Where  $L_R$  is some smooth function on  $U_R = \{x \in M | R - \frac{r}{n(n-1)}G \neq 0 \text{ at } x\}$ ,  $G$  is the  $(0, 4)$ -tensor defined by  $G(X_1, X_2, X_3, X_4) = g((X_1 \wedge X_2)X_3, X_4)$  and  $(X_1 \wedge X_2)X_3$  is the endomorphism and it is defined as,

$$(X_1 \wedge X_2)X_3 = g(X_2, X_3)X_1 - g(X_1, X_3)X_2. \quad (27)$$

**Definition 4.** In a Riemannian manifold  $M$ , if  $R \cdot S$  and  $Q(g, S)$  are linearly dependent then  $M$  is called a Ricci pseudo symmetric manifold and this condition is given by

$$R \cdot S = f_s Q(g, S), \quad (28)$$

where  $U_S = \{x \in M : S - \frac{k}{n} \neq 0 \text{ at } x\}$  and  $f_s$  is a function defined on  $U_S$ .

**Theorem 5.** Let  $(2n + 1)$ -dimensional Ricci pseudosymmetric generalized  $(k, \mu)$ -space form is Ricci semisymmetric, provided  $f_s \neq f_1 - f_3$  and  $f_4 = f_6$ .

*Proof.* Now equation (28) can be written as

$$(R(X, Y)S)(Z, W) = -f_s\{S((X \wedge_g Y)Z, W) + S(Z, (X \wedge_g Y)W)\}, \quad (29)$$

where the endomorphism  $(X \wedge_g Y)Z = g(Y, Z)X - g(X, Z)Y$ . Now (29) yields

$$-S(R(X, Y)Z, W) - S(Z, R(X, Y)W) = -f_s\{S(Y, W)g(X, Z) - S(X, W)g(Y, Z) + S(Z, Y)g(X, W) - S(Z, X)g(Y, W)\}. \quad (30)$$

Putting  $X = Z = \xi$  in (30), we get

$$-S(R(\xi, Y)\xi, W) - S(\xi, R(\xi, Y)W) = -f_s\{S(Y, W)g(\xi, \xi) - S(\xi, W)g(W, \xi) + S(\xi, Y)g(\xi, W) - S(\xi, \xi)g(Y, W)\}. \quad (31)$$

Using (1), (6) and (7) in the above equation, we have

$$((f_1 - f_3) - f_s)[S(Y, W) - 2n(f_1 - f_3)g(Y, W)] + (f_4 - f_6)[S(hY, W) - 2n(f_1 - f_3)g(hY, W)] = 0. \quad (32)$$

Hence proof. Further, from (32) We can state the following statement.

**Theorem 6.** A  $(2n + 1)$ -dimensional Ricci pseudosymmetric generalized  $(k, \mu)$ -space form is Ricci pseudosymmetric generalized Sasakian space form if and only if  $(f_4 - f_6) = 0$ .

In [22] Tripathi et.al., introduced  $\tau$ -curvature tensor and which is the generalization of conformal, concircular, projective etc. This curvature tensor was studied on  $K$ -contact, Sasakian and  $(k, \mu)$ -contact metric manifolds by the authors in [21, 22]. The  $\tau$ -curvature tensor is defined by [22]

$$\begin{aligned} \tau(X, Y)Z &= a_0R(X, Y)Z + a_1S(Y, Z)X + a_2S(X, Z)Y + a_3S(X, Y)Z + a_4g(Y, Z)QX \\ &\quad + a_5g(X, Z)QY + a_6g(X, Y)QZ + a_7r\{g(Y, Z)X - g(X, Z)Y\}. \end{aligned} \quad (33)$$

**Definition 5.** A  $(2n + 1)$ -dimensional generalized  $(k, \mu)$ -space form is said to be  $\tau$ -flat if it satisfies

$$\tau(X, Y)Z = 0. \quad (34)$$

**Theorem 7.** A  $\tau$ -flat  $(2n + 1)$ -dimensional generalized  $(k, \mu)$ -space form is an  $\eta$ -Einstein manifold.

*Proof.* Consider a  $\tau$ -flat  $(2n + 1)$ -dimensional generalized  $(k, \mu)$ -space form. Then from (34) we have

$$\begin{aligned} a_0 g(R(X, Y)Z, W) &= -a_1S(Y, Z)g(X, W) - a_2S(X, Z)g(Y, W) - a_3S(X, Y)g(Z, W) - a_4g(Y, Z)S(X, W) \\ &\quad - a_5g(X, Z)S(Y, W) - a_6g(X, Y)S(Z, W) - a_7r\{g(Y, Z)g(X, W) - g(X, Z)g(Y, W)\}. \end{aligned} \quad (35)$$

Putting  $X = W = \xi$  in the above equation and using (6) and (1), it follows that

$$S(Y, Z) = Ag(Y, Z) + B\eta(Y)\eta(Z),$$

where

$$A = -\left\{ \frac{(2n-1)f_4 - f_6(a_4 2n(f_1 - f_3) + a_7r) - a_0(f_4 - f_6)(2nf_1 + 3f_2 - f_3)}{a_0(f_4 - f_6) - a_1((2n-1)f_4 - f_6)} \right\}$$

and

$$B = -\left\{ \frac{((2n-1)f_4 - f_6)(-a_0 + 2n(a_2 + a_3 + a_5 + a_6))(f_1 - f_3) + a_7r + (3f_2 + (2n-1)f_3)}{a_0(f_4 - f_6) - a_1((2n-1)f_4 - f_6)} \right\}.$$

This complete the proof of the theorem.

**Definition 6.** A Riemaniann manifold  $(M, g)$  satisfying the condition [21, 22]

$$\phi^2(\tau(\phi X, \phi Y)\phi Z) = 0, \quad (36)$$

is called  $\phi$ - $\tau$  flat.

Suppose that the generalized  $(k, \mu)$ -space form  $M^{2n+1}(f_1, \dots, f_6)$  is  $\phi$ - $\tau$ -flat. Then

$$g(\tau(\phi X, \phi Y)\phi Z, \phi W) = 0, \quad (37)$$

for all vector fields  $X, Y, Z$  and  $W$ .

**Theorem 8.** A  $\phi$ - $\tau$  flat generalized  $(k, \mu)$ -space form  $M^{2n+1}(f_1, \dots, f_6)$  is an  $\eta$ -Einstein manifold.

*Proof.* We know that  $M^{2n+1}(f_1, \dots, f_6)$  is  $\phi$ - $\tau$  flat, (33) can be written as

$$\begin{aligned} a_0 g(R(\phi X, \phi Y)\phi Z, \phi W) &= -a_1 S(\phi Y, \phi Z)g(\phi X, \phi W) - a_2 S(\phi X, \phi Y)g(\phi Z, \phi W) \\ &\quad - a_3 S(\phi X, \phi Y)g(\phi Z, \phi W) - a_4 g(\phi Y, \phi Z)S(\phi X, \phi W) \\ &\quad - a_5 g(\phi X, \phi Z)S(\phi Y, \phi W) - a_6 g(\phi X, \phi Y)S(\phi Z, \phi W) \\ &\quad - a_7 r [g(\phi Y, \phi Z)g(\phi X, \phi W) - g(\phi X, \phi Z)g(\phi Y, \phi W)]. \end{aligned} \tag{38}$$

Let  $\{e_1, \dots, e_{n-1}, \xi\}$  be a local orthonormal basis of vector fields in  $M^{2n+1}(f_1, \dots, f_6)$  and using the fact that  $\{\phi e_1, \dots, \phi e_{n-1}, \xi\}$  is also a local orthonormal basis, if we put  $Y = Z = e_i$  in (38) and sum up with respect to  $i$ , then we get

$$\begin{aligned} a_0 \sum_{i=1}^{2n} g(R(\phi X, \phi e_i)\phi e_i, \phi W) &= -a_1 \sum_{i=1}^{2n} S(\phi e_i, \phi e_i)g(\phi X, \phi W) - a_2 \sum_{i=1}^{2n} S(\phi X, \phi e_i)g(\phi e_i, \phi W) \\ &\quad - a_3 \sum_{i=1}^{2n} S(\phi X, \phi e_i)g(\phi e_i, \phi W) - a_4 \sum_{i=1}^{2n} g(\phi e_i, \phi e_i)S(\phi X, \phi W) \\ &\quad - a_5 \sum_{i=1}^{2n} g(\phi X, \phi e_i)S(\phi e_i, \phi W) - a_6 \sum_{i=1}^{2n} g(\phi X, \phi e_i)S(\phi e_i, \phi W) \\ &\quad - a_7 r \sum_{i=1}^{2n} [g(\phi e_i, \phi e_i)g(\phi X, \phi W) - g(\phi X, \phi e_i)g(\phi e_i, \phi W)]. \end{aligned} \tag{39}$$

It can be easily verified that,

$$\sum_{i=1}^{2n} g(R(X, \phi e_i)\phi e_i, W) = S(\phi X, \phi W) + g(\phi X, \phi W), \tag{40}$$

$$\sum_{i=1}^{2n} g(e_i, e_i) = \sum_{i=1}^{2n} g(\phi e_i, \phi e_i) = 2n, \tag{41}$$

$$\sum_{i=1}^{2n} S(\phi e_i, \phi e_i) = r - 2n(f_1 - f_3), \tag{42}$$

$$\sum_{i=1}^{2n} g(X, \phi e_i)g(\phi e_i, W) = g(\phi X, \phi W). \tag{43}$$

Using (3), (8) and (40)-(43) in (39) we get

$$S(X, W) = Ag(X, W) + B\eta(X)\eta(W).$$

where

$$A = -\left\{ \frac{a_0 + a_4(r - 2n(f_1 - f_3)) + a_7r(2n - 1)}{a_0 + 2na_1 + a_2 + a_3 + a_5 + a_6} \right\}$$

and

$$B = \left\{ \frac{a_0 + a_4(r - 2n(f_1 - f_3)) + a_7r(2n - 1) + 2n(f_1 - f_3)(a_0 + 2na_1 + a_2 + a_3 + a_5 + a_6)}{a_0 + 2na_1 + a_2 + a_3 + a_5 + a_6} \right\}.$$

Hence the result.

## Acknowledgement

The first authors is thankful to University Grants Commission, New Delhi, India for financial support in the form of National Fellowship for Higher Education (F1-17.1/2016-17/NFST-2015-17-ST-KAR-3079/(SA-III/Website)).

## Competing interests

The authors declare that they have no competing interests.

## Author's contributions

All authors have contributed to all parts of the article. All authors read and approved the final manuscript.

## References

- [1] P. Alegre, D. E. Blair and A. Carriazo, *Generalized Sasakian Space forms*, Israel J. Math., 141 (2004), 157-183.
- [2] P. Alegre and A. Carriazo, *Structures on generalized Sasakian space forms*, Differential Geometry and its Applications, 26 (2008), 656-666.
- [3] R. Al-Ghefari, F. R. Al-Solamy and M. H. Shahid, *Contact CR-warped product submanifolds in generalized Sasakian space forms*, Balkan J. Geom. Appl., 11 (2) (2006), 1-10.
- [4] C. S. Bagewadi and M. C. Bharathi, *A study on hypersurface of complex space form*, Acta Et Commentationes Universitatis Tartuensis De Mathematica, 17 (1) (2013), 65-70.
- [5] D. E. Blair, *Contact manifolds in Riemannian geometry*, Lecture Notes in Mathematics, 509 (1976), Springer-Verlag, Berlin.
- [6] D. E. Blair, *Riemannian Geometry of Contact and Symplectic Manifolds*, Birkhäuser Boston, 2002.
- [7] E. Cartan, *Leçons sur la géométrie des espaces de Riemann*, Gauthier-Villars Paris, (1963).
- [8] A. Carriazo and V. M. Molina, *Generalized  $(K, \mu)$  space forms and  $D_\alpha$  homothetic deformations*, Balkan Journal of Geometry and Its Applications, 16 (1) (2011), 37-47.
- [9] A. Carriazo, V. M. Molina and M. M. Tripathi, *Generalized  $(K, \mu)$  space forms*, Mediterranean Journal of Mathematics, 10 (2013), 475-496.
- [10] M. C. Chaki, *On pseudo symmetric manifolds*, Ann. St. Univ. Al I Cuza Iasi, 33 (1987).
- [11] B. Y. Chen, *Geometry of submanifolds*, Marcel Dekker. Ine. New York, (1973).
- [12] U. C. De and P. Majhi, *Certain Curvature properties of generalized Sasakian space forms*, Facta Universitatis (NIS)Ser. Math. Inform., 27 (3) (2012), 271-282.
- [13] U. C. De and S. Mallick, *On the existence of generalized quasi Einstein manifolds*, Archivum Mathematicum (Brno), 47 (2011), 279-291.
- [14] R. Deszcz, *On pseudosymmetric spaces*, Bull. Soc. Math. Belg. Ser. A, 44 (1992), 1-34.
- [15] F. Gherib and M. Belkhef, *Parallel submanifolds of generalized sasakian space forms*, Bulletin of the Transilvania University of Brasov Series III, 2 (51) (2009), 185-192.
- [16] A. Ghosh, *Killing Vector Fields and Twistor Forms on Generalized Sasakian Space Forms*, Mediterr. J. Math., 10 (2013), 1051-1065.
- [17] T. Koufogiorgos, *Contact Riemannian manifolds with constant  $\phi$ -sectional curvature*, Tokyo J. Math., 20 (1) (1997), 55-67.
- [18] F. Malek and V. Nejadakbary, *Warped product submanifold in generalized sasakian space form*, Acta Mathematica Academiae Paedagogicae Nyiregyhaziensis., 27 (2011), 325-338.
- [19] A. Olteanu, *Legendrian warped product submanifolds in generalized sasakian space forms*, Acta Mathematica Academiae Paedagogicae Nyiregyhaziensis., 25 (2009), 137-144.
- [20] S. Tano, *Ricci curvatures of contact Riemannian manifolds*, Tohoku Math.J., 40 (1988), 441-448.



- [21] M. M. Tripathi and P. Gupta , *On  $\tau$ -curvature tensor in K-contact and Sasakian manifolds*, International Electronic Journal of Geometry, 4 (2011), 32-47.
- [22] M. M. Tripathi and P. Gupta,  *$\tau$ -curvature tensor on a semi-Riemannian manifold*, J. Adv. Math. Stud., 4 (1) (2011), 117-129.
- [23] S. Sular and C. Ozgur, *On some submanifolds of Kenmotsu manifolds*, Chaos, Solitons and Fractals, 42 (2009), 1990-1995.
- [24] Z. Szabo, *Structure theorems on Riemannian spaces satisfying  $R(X,Y) \cdot R = 0$ . I. the local version*, J. Differ Geom., 17 (1982), 531-582.
- [25] S. Sular and C. Ozgura, *Contact CR-warped product submanifolds in generalized Sasakian space forms*, Int. Electron. J. Geom., 4 (1) (2011), 32-47.
- [26] Venkatesha and B. Shanmukha, *M-Projective Curvature Tensor on Lorentzian  $\alpha$ -Sasakian Manifolds*, Global Journal of Pure and Applied Mathematics, 13 (7) (2017), 2849-2858.
- [27] Venkatesha and B. Sumangala, *on M-projective curvature tensor of Generalised Sasakian space form*, Acta Math., Univ. Comenianae 2 (2013), 209-217.